

# MASTIVAC

Reg. SAGARPA.: B-0029-001

Bacterin for the prevention of bovine mastitis.



## Formula

Each mL contains inactivated cultures of:

<i>Staphylococcus aureus</i> .....	≥12 X 10 <sup>8</sup> UFC
<i>Streptococcus agalactiae</i> .....	≥9 X 10 <sup>8</sup> UFC
<i>Streptococcus dysgalactiae</i> .....	≥3 X 10 <sup>8</sup> UFC
<i>Streptococcus uberis</i> .....	≥3 X 10 <sup>8</sup> UFC
<i>Streptococcus pyogenes</i> .....	≥3 X 10 <sup>8</sup> UFC
<i>Escherichia coli</i> .....	≥18 X 10 <sup>8</sup> UFC
<i>Corynebacterium pyogenes</i> .....	≥12 X 10 <sup>8</sup> UFC
Adjuvant 20% of final volume	
Diluent: Water injection	



## Indications

Prevention of bovine mastitis.



## Administration and dose

Administer two injections of 5 mL each one with interval of 15 days between each injection.  
Primo-vaccination: Two months before calving.  
Re-vaccination: Annual.  
Subcutaneous in the neck or behind the back.



## Instructions and warnings

Keep refrigerated between 2°C and 8°C. Keep in a cool place and protected from light. Bottles used and any leftovers should be removed in a safety way for environment and according to local and national regulations.

Use in:



Presentations:

20 doses bottle (100 mL)



VETERINARY USE  
FOR THE USE OF THE VETERINARIAN  
CONSULT THE VETERINARIAN